

Extraction of β -glucan from oats and development of functional β -glucan enriched beverage

Jasia Nissar¹, Tehmeena Ahad²

^{1,2}Division of Food Science and Technology,

Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir

ABSTRACT

Oats (*Avena Sativa*), contain β -glucan as a source of soluble dietary fiber. The oat flour was analyzed for its chemical composition. The oat flour possessed 11.6% total dietary fibre and 4.8% β -glucan content, 6.2% fat, 16.1% protein and 66.1% carbohydrates. The beverage was prepared by incorporating β -glucan at 0, 0.4, 0.8 and 1.0% levels. The L^* -value of beverage increased, while b^* -value decreased progressively by incorporation of oat β -glucan. The viscosity and acidity of the beverage also improved significantly with β -glucan level. Sensory evaluation revealed that the beverage containing 0.2 and 0.4% β -glucan showed similar response as that of control. Overall the incorporation of β -glucan upto 0.8% did not significantly affect sensory parameters of flavor, color and acceptability.

Key Words: β -glucan; Extraction; Functional beverage; Oats.

I. INTRODUCTION

Oats are gaining importance as an ingredient in the production of functional foods due to its higher content of bioactive compounds. It possesses high amount of dietary fiber (DF) with high proportion of soluble viscous components, offering more suitability among cereal grains in the human diet (1). The oats contain substantially higher amounts of functional ingredient β -glucan. The use of β -glucan extracted from oats as human food due to its positive role in human health has received a growing attention. The cell wall of barley and oat contains β -glucan, a non-starch polysaccharide composed of β -(1-4)-linked glucose units separated every two to three units by a single β -(1-3)-linked glucose and referred to as a mixed linkage β -glucan (2). β -glucan delays glucose absorption and regulates the level of blood glucose (3). The viscous nature of β -glucan physically slows glucose absorption in the gut. This property of β -glucan may be useful in the formulation of food products targeting management of diabetes. Functional foods including functional beverages are important for their role in health promotion and disease prevention. The oats can be used to enhance the flavor, texture, appearance and nutritional composition in a variety of functional foods, including hot cereals, cookies, crackers, breads, tortillas, granola bars, fruit-filled cereal bars, extruded snacks and pastas and development of different beverages (4). Growing interest in new functional foods with special characteristics and health benefits has led to the development of new functional beverages. These beverages may enrich diet and improve human health, because of its ease of consumption along with a usual meal. oat β -glucan assume to be well suited for such a functional

application, being capable of imparting a smooth mouth feel to beverage products and providing an excellent source of soluble dietary fiber. β -glucan gum with similar functional properties, could potentially serve as an alternative to traditional beverage thickeners such as alginates, pectin, xanthan and carboxymethylcellulose (5). The present study was planned to extract the β -glucan from oats and utilize it for the development of functional beverage.

II. MATERIALS AND METHODS

Preparation and analysis of oat flour. oat flour was prepared by grinding oat grains through UDY cyclone mill (mesh size 20 mm). The flour was analyzed for proximate composition i.e., moisture, crude fat, crude protein, ash, crude fiber and nitrogen free extract (NFE) by following methods of AACC (2000).

Extraction and purification of β -glucan. β -glucan gum was extracted from oatflour by following with the method of Wood *et al.* (1978) with some modifications. The oat flour (50 g) was suspended in 500 mL water, pH was adjusted to 10 with Na₂CO₃ (20%, v/w) and stirred vigorously for 30 min at a temperature of 45°C. The mixture was centrifuged at 15000 x g at 5°C for 15 min. The supernatant was adjusted to pH 4.5 with 2M HCL and centrifuged again to separate precipitated protein, which was discarded. The β -glucan was precipitated by adding of an equal volume of ethanol (99.9%) to the supernatant with slow stirring. The precipitate was recovered by centrifugation at 3300 x g for 10 min allowed to settle overnight at 5°C and dried in a vacuum drier. The extracted β -glucan was stored in high density polyethylene (HDPE) bags at 28 oC for further studies.

Utilization of β -glucan.

The purified β -glucan was utilized in different formulations for the preparation of functional beverages, as given in Table I.

Preparation and evaluation of β -glucan beverage. The β - glucan beverage was prepared following the formulation of Temelli *et al.* (2004) with some modifications . The functional beverage samples were then organoleptically evaluated for sensory parameters such as color, flavor, sweetness, sourness and overall acceptability by a panel of five judges. The nine point hedonic scale was employed for the evaluation of samples.

Physicochemical evaluation of β -glucan beverage. The color values of β -glucan beverage samples were measured according to method of Yu *et al.* (2003) by using the L* a* b* color space (CIELAB Space) with Color Tech-PCM (USA). The acidity of beverage samples was determined by following the method given in AOAC (2000). The viscosity of functional beverages was measured by following the procedure of AACC (2000) through viscometer (Rion Tech., USA).

Statistical analysis. The data were subjected to analysis of variance (ANOVA) using CoStat-2003 software following the method as described by Steel *et al.* (1997). The Duncun Multiple Range (DMR) was used to determine the level of significance between samples.

III.RESULTS AND DISCUSSION

Chemical composition of oat flour. The oat flour contained 16.1%, 6.2%, 11.6%, 2.22% and 66.1% crude protein, crude fat, dietary fiber, ash and carbohydrate, respectively (Table II), which corroborated the earlier findings for Canadian varieties by (11).

Physicochemical evaluation of β -glucan beverages. The color values of beverage samples indicated that the L*-value (color index) of functional beverages increased significantly as the level of β -glucan increased in the formulation of different beverages. The beverage containing 1.0% β -glucan showed the highest L*-value (23.31) while control beverage (without β -glucan), which got L*-value 20.82. However the beverage containing 1% β -glucan gave the lowest a*-value (1.29) and the highest a*-value (1.54) was given by control beverage (without β -glucan). Similarly, b*-value was significantly affected by treatments. The control beverage containing 0.2% pectin possessed the highest b*-value (9.89) followed by the beverage containing 1% β -glucan. and significantly the lowest b*-value was recorded in the beverage with 0.2% β -glucan. The present study indicated that incorporation of β -glucan resulted in improvement of beverages color as compared to the control beverage, which was prepared by the addition of 0.2% pectin without addition of β -glucan. A small amount of precipitate was visible at the bottom of the β -glucan beverage, which is due to insoluble protein and fiber components present in the β -glucan at low levels. Thus the precipitation of this material in case of β -glucan supplemented beverage might be a cause of higher L*-value for these treatments of beverage. In the present study a*-value decreased significantly by increasing the level of β -glucan in the beverages, which indicated that increased β -glucan concentration resulted in a less reddish product as compared to the control beverage. The viscosity of beverages improved significantly due to the incorporation of β -glucan in beverages. The highest viscosity (26.32 mPa-s) was found in beverages containing 1% β -glucan followed by that containing 0.8% β -glucan (Table III). The addition of β -glucan to water also results in the formation of a viscous hydrocolloid solution (12), which might be one of the reasons towards increase in the viscosity of beverages. The polysaccharide's hydroxyl groups are available to form hydrogen bonds with water, which makes the polymer water-soluble. Total acidity varied significantly as a function of treatment. The variation in acidity in the present study was due to the degradation of sucrose, high fructose corn syrup and β -glucan by the action of microorganisms, which causes production of acids in beverages (13).

Sensory evaluation of β -glucan beverages. The scores assigned to the sensoric attributes of 0.2% of β -glucan containing beverages revealed that these beverage got significantly the higher color scores (7.20) followed by the control beverage (0.2% pectin). Similar trend was observed in case of sweetness and sourness, however the scores assigned to beverage prepared by the incorporation of 0.4% β -glucan got the highest score for flavor (Table IV). Contrarily the control treatment (0.2% pectin) got the top position regarding overall acceptability of beverage, followed by beverage samples prepared by incorporation of 0.4% β -glucan; with no significant differences in both treatments. β -glucan's ability to increase viscosity upon addition to water makes it an

excellent thickener for beverage applications. These characteristics are more appealing to the panelists for making decision about the overall acceptability of beverages. The results further indicated that in the beverages fortified with polysaccharides like β -glucan the quality characteristics of the beverages varies nonsignificantly but it was also revealed that the incorporation should be not more than 0.4% of β -glucan. The further increase in β -glucan level thickens the beverage and higher consistency not appeals the consumer regarding sensory prospective of beverage.

IV. CONCLUSION

Incorporation of β -glucan has an effect on physicochemical characteristics and sensoric of beverage. The beverage improved regarding most of the physicochemical and sensory characteristics of the beverage. The acidity and viscosity of beverage improved linearly as the incorporation level of β -glucan increased in beverage formulation. Similarly the consumers like more viscous beverage, which was prepared by incorporation of β -glucan. However the beverages containing lesser than 0.8% β -glucan were the least acceptable by the panelists. Further research is needed to know the thermal stability of β -glucan and its behavior with other food ingredients in beverages application to make stable foods.

Table I. Incorporation level of β -glucan

Treatment	β-glucan incorporation
1	0control (0.2% pectin)
2	0.4%
3	0.8%
4	1%

Table II. Chemical composition of oat flour

Component	% (dry basis)
Crude protein	16.1
Crude fat	6.2
Dietary fiber	11.6
Ash	2.22
Carbohydrates	66.1

Table III. Effect of β -glucan incorporation on physico-chemical properties of beverage

% β -glucan	Acidity	Viscosity	L*	a*	b*
0	1.48±0.02	4.42±0.06	20.85	1.54	9.89
0.2	1.50±0.05	8.01±0.19	20.89	1.48	9.61
0.4	1.53±0.04	16.48±0.43	21.20	1.41	9.43
0.8	1.52±0.04	20.88±0.84	21.93	1.32	9.26
1	1.53±0.04	26.32±0.75	22.31	1.29	8.86

Table IV. Effect of β -glucan incorporation on overall acceptability of beverage

% β -glucan	Overall acceptability
0	7.18
0.2	7.20
0.4	5.46
0.8	5.28
1	5.28

REFERENCES

- [1.] Bjorck, I., A.C. Eliasson, A. Drews, M. Gudmundsson and R. Karlsson, 1990. Some nutritional properties of starch and dietary fiber in barley genotypes containing different levels of amylose. *Cereal Chem.*, 67: 327–336
- [2.] Carpita, N.C., 1996. Structure and biogenesis of cell walls of grasses. *Annl. Rev. Plant Physiol. Plant Mol. Biol.*, 47: 445–476
- [3.] Wood, P.J., J.T. Braaten, F.W. Scott, K.D. Riedel, M.S. Wolynetz, M.W. Collins, 1994. Effect of dose and modification of viscous properties of oat gum on plasma glucose and insulin following an oral glucose load. *British J. Nutri.*, 72: 731–743
- [4.] Arndt, E.A., 2006. Whole-grain barley for today's health and wellness needs. *ConAgra Foods, Inc. Omaha, NE.*, 51: 20–22.
- [5.] Giese, J.H., 1992. Hitting the spot: Beverages and beverage technology. *Food Technol.*, 46: 70–72
- [6.] AACC, 2000. *Approved Methods of American Association of Cereal Chemists*. The American Association of Cereal Chemists, Inc. St. Paul, Minnesota, USA
- [7.] Wood, P.J., I.R. Siddiqui and D. Paton, 1978. Extraction of High-Viscosity Gums from Oats. 1978. *Cereal Chem.*, 55: 1038–1049

- [8.] Temelli, F., C.B. Bansema and K.S. Stobbe, 2004. Development of an orange flavored barley β -glucan beverage. *CerealChem.*, 81: 499–503
- [9.] Yu, L., J. Perret, M. Harris, J. Wilson and S. Haley, 2003. Antioxidant properties of bran extracts from Akron wheat grown at different locations. *J. Agric. Food Chem.*, 51: 1566–1570
- [10.] Steel, R.G.D., J.H. Torrie and D.A. Dickey, 1997. *Principles and Procedures of Statistics: A Biometrical Approach*, (3rd edition). McGraw Hill Book Co. Inc., New York, USA
- [11.] LI, J.H., T. Vasanthan, B. Rossnagel and R. Hoover, 2004. Starch from hull-less barley: I. Granule morphology, composition and amylopectin structure. *Food Chem.*, 74: 395–405
- [12.] Dawkins, N.L. and I.A. Nnanna, 1993. Studies on oat gum [(1 \rightarrow 3, 1 \rightarrow 4)- β -D-glucan]: Composition, molecular weight estimation and rheological properties. *Food Hydrocol.*, 9:
- [13.] Renuka, A.B., S.G. Kulkarnib, P. Vijayanandb and S.G. Prapulla, 2009. Fructooligosaccharide fortification of selected fruit juice beverages: Effect on the quality characteristics. *Food Sci. Technol.*, 1–3