

CHIA SEED (*SALVIA HISPANICA L.*) – A NEW AGE FUNCTIONAL FOOD

Dr Sukhneet Suri¹, Dr Santosh Jain Passi², Jyoti Goyat³

¹Assistant Professor, Department of Food Technology,

Vivekananda College, University of Delhi, (India)

²Former Director, Institute of Home Economics, University of Delhi, (India)

Public Health Nutrition Consultant, Hon. Professor-PHN, Amity University, Greater Noida, (India)

³Ph.D. Research Scholar, Amity University, Greater Noida, (India)

ABSTRACT

A review has been made on chia seeds (*Salvia hispanica L.*), a non-conventional seed which is increasingly being recognized as a novel food and is receiving scientific attention. Chia is an annual herb. It is a good source of omega-3/omega-6 fatty acids, soluble dietary fiber and contains appreciable amount of proteins and phytochemicals. It thus has nutritional attributes which support the prevention of several non-communicable diseases such as obesity, hypertension, cardio-vascular disease (CVD's), cancer and diabetes. This review presents an overall yet comprehensive view of the present state of knowledge regarding the history, structure, nutritional composition and physicochemical properties of Chia. It critically brings forth the effects of consumption of Chia seeds on human physiology.

In this systematic literature review (with no meta-analysis) scientific research and review papers (N=50) were collected via several electronic databases such as Medline (Pub-Med Version), NLIST (Programme of INF LBNET) and DELNET (Developing library network) under the keywords "Chia seeds", "Salvia hispanica", "Hypertension", "Origin", "Hyperglycemia", "Cardiovascular risk", "Triglycerides", "Omega-3 fatty acids", "Nutraceutical", "Dietary fiber", "Fatty acids", "Physicochemical properties", "Functions", "Viscosity", "Mucilage", "Antioxidants", "Phenolic compounds", "Viscosity", "Morphology", "Applications", "Gelling", "Plasma levels", "Clinical trials", "Insulin Resistance", "Adiposity".

Keywords: Antioxidant, Chia Seeds, Non- Communicable Disease, Novel food, Omega-3 fatty acid.

I. INTRODUCTION

Chia seeds or *Salvia hispanica L.* belong to the Lamiaceae (Mint) family. Chia is an annual herb, native to Mexico and Northern Guatemala which blooms in summer season [1, 2]. *Salvia hispanica* was used as food as early as in 3500 BC and gained importance as a crop between 1500 BC and 900 BC in central Mexico. It was used as a staple crop by Aztecs and was also offered to Gods in religious ceremonies. This ancestral seed has been known for its medicinal and nutritional properties from ancient time because of its high content of omega-3 fatty acids. While chia seeds were used as food, chia oil was used in cosmetics and paintings by Aztecs [3]. Nowadays chia seeds are commercially grown in Mexico, Bolivia, Argentina, Ecuador and Guatemala [4].

II. STRUCTURE AND COMPOSITION

Chia seeds are generally small, flat and oval shaped. The seeds are 2.0 - 2.5 mm long, 1.2 - 1.5 mm wide and 0.8 – 1.0 mm in thickness. The chia plant is sensitive to day light and produces black and white seeds. Black colored chia seeds are more common. The black and white colour seeds are slightly different from each other. White seeds are larger, thicker and broader than the black seeds. The average moisture content of black and white seeds is 7.2 per cent and 6.6 per cent respectively. The oil yield of white seeds and black seeds is about 33.8 per cent and 32.7 per cent respectively. Protein and fatty acid composition of both the seeds varies significantly [1, 3].

Chia seeds are a good source of fatty acids, dietary fiber, protein, minerals and antioxidants. Chia seeds can be used as whole, milled or grounded. Oil extracted from chia seeds can also be used in food [2, 5].

III. NUTRITIONAL COMPOSITION OF CHIA SEEDS

The nutrient composition of chia seeds have been assessed and studied by several researchers and organizations. The seeds are considered to be of nutritional significance in view of their fatty acid composition as well as the fiber and protein content. TABLE 1 gives the detailed macro-nutrient composition of chia seeds.

TABLE 1: Comparison of the meannutritional composition of chia seeds with other grains/seeds.

Grains/Seeds	Chia	Flax	Quinoa	Sunflower	Mustard	Guar	Oats*
Nutrients Amount per 100g	Seeds	seeds	seeds	seeds	seeds	Gum seeds	
Energy(Kcal)	486	534	368	584	508	332	374
Protein(g)	16.54	18.29	14.12	20.78	26.08	4.60	13.6
Fat(g)	30.74	42.16	6.07	51.46	36.24	0.50	7.6
Carbohydrates (by difference; g)	42.12	28.88	64.16	20.0	28.09	77.3	66.27
Moisture (g)	5.80	6.96	13.28	4.73	5.27	15	8.22

Source: USDA National Nutrient Database for Standard Reference Release 28, 2011[6] and *NIN, Nutritive value of Indian Foods [7]

Different ecosystems have variable significant effects on the nutrient composition of *Salvia hispanica* especially its protein and oil content and also the fatty acid composition. The environmental factors which have been found to influence composition of chia seeds include temperature, light, soil composition and type/variety [8]. Analysis of chia seeds grown commercially in three different ecosystems namely Argentina (T₁ - Semi Arid Chaco Ecosystem), Bolivia (T₂ - Sub Humid Chaco Ecosystem) and Ecuador (T₃ - Inter- Andean valley) has indicated that the protein content decreases as altitude increases. Chia seeds grown in the sub humid chaco ecosystem (T₂) have been found to contain 61 per cent more protein than other two ecosystems i.e. the semi-arid chaco and the inter-andean valley ecosystem. Oil content does not show any significant difference among the different ecosystems but its composition varies. Chia seeds cultivated in ecosystem T₃ contain higher concentration of

alpha linolenic acid (56.9per cent to 64.8per cent) which is the main constituent of oil. Since chia seeds are the richest source of omega-3 fatty acids among plant foods and are a good source of fiber and phytochemicals, they can be considered as functional food[9].

3.1 Protein

The average protein content varies from 15per cent to 23per cent according to the location where the seeds have been grown [10]. Chia seeds (per 100g) contain higher amount of protein (16.54 g) as compared to other grains like wheat (11.8 g), oats (13.6 g), barley (11.5 g), rice (6.8 g) and corn (11.1 g) [3, 6,7]. The protein content of defatted chia flour as assessed by dry fractionation is about 446.2g/kg of proteins [11]. TABLE 2 gives the detailed amino acid composition of chia seeds.

TABLE 2: Average amino-acid composition of chia seeds

Amino acid (g per 100g)		Amino acid (g per 100g)	
Arginine	2.143	Phenylalanine	1.016
Glutamic acid	3.500	Tyrosine	0.536
Threonine	0.709	Histidine	0.531
Tryptophan	0.436	Valine	0.950
Isoleucine	0.801	Alanine	1.044
Leucine	1.371	Glycine	0.943
Methionine	0.588	Aspartic acid	1.689
Lysine	0.970	Proline	0.776
Cystine	0.407	Serine	1.049

Source: USDA National Nutrient Database for Standard Reference Release 28, 2011 [6].

A comparison of the amino acid profile of chia seeds with other seeds is given in fig. 1 [6,12].




Figure 1: Amino acid composition of chia seeds and other seeds/grains – a comparison

3.2 Dietary fiber

Fiber is one of the important components of healthy diet. Intake of adequate amount of dietary fiber is associated with the prevention of cardiovascular diseases like stroke, myocardial infarction, vascular diseases, obesity, hypertension, hyperglycemia, and hyperlipidemia. Dietary fibers cannot be digested and absorbed by the small intestine but get fermented in the large intestine. On the basis of their physio-chemical properties and functions, dietary fibers are classified in two forms- insoluble fibers which exhibit bulking action and soluble

dietary fibers which get fermented partially/completely in the colon [13, 14]. The total dietary fiber content of chia seeds ranges from 36 to 40 g per 100g which is much higher than that present in several grains, vegetables and fruits such as corns, carrot, spinach, banana, pear, apple, kiwi [5, 11]. Insoluble and soluble dietary fiber varies from 23 to 46 per cent and 2.5 to 7.1 per cent respectively [15]. Chia contains about 5 per cent mucilage which can also act as soluble fiber [16]. The insoluble dietary fibre of chia is capable of retaining water several times of its weight during hydration and thus provides bulk and prolongs the gastro-intestinal transit time. Increased gastro-intestinal time is directly related to gradual increase in post-prandial blood glucose levels and decrease in insulin resistance over a period of time [16].

TABLE 3: Dietary fiber content of different foods

Food stuff	Chia seeds	Flax seeds	Quinoa seeds	Guar gum seeds
Amount per 100g	34.4g	27.3g	7.0g	77.3g

Source: USDA National Nutrient Database for Standard Reference Release 28, 2011 [6].

3.3 Total Fats

Salvia hispanica contain on an average 30.74 per cent of total lipids [6]. Several researchers have assessed the fatty acid composition of chia seeds and variable results have been obtained. For example, in a study carried out by Marineli et al (2014) alpha linolenic fatty acid was found to be in most abundance (62.80g/100g) followed by linoleic acid (18.23g/100g), palmitic acid (7.07g/100g), oleic acid (7.04g/100g) and stearic acid (3.36g/100g) [17]. This fatty acid composition however varies from that given by United States Department of Agriculture. Refer TABLE 4 for the fatty acid composition given by USDA(2011). Chia seeds contain on an average, 40 per cent of their total weight as oil. Their oil contains nearly 60 per cent omega-3 unsaturated fatty acids [5]. Omega-3 unsaturated fatty acids are helpful in the prevention and management/treatment of hyperlipidemia, hyperglycemia and hypertension [2, 18, 19]. Chia seeds contain higher concentration of omega-3 fatty acid (63.8per cent) as compared to flaxseeds (57.5per cent) and menhaden fish oil (1.5per cent) [20].

TABLE 4: Average fatty acid composition of chia seeds

Total Saturated fatty acids (g/ 100g)	3.330g	Total Monounsaturated fatty acids (g/ 100g)	2.309g	Total Polyunsaturated fatty acids g/100g)	23.665g
14:0	0.030g	14:1	0.030g	18:2 n-6	5.835g
15:0	0.030g	16:1	0.029g	18:3	17.830g
16:0	2.170g	17:1	0.000g	Trans fatty acids	0.140g
17:0	0.063g	18:1	2.203g	Cholesterol	0mg
18:0	0.912g	20:1	0.046g		
20:0	0.093g				
22:0	0.032g				

Source: USDA National Nutrient Database for Standard Reference Release 28, 2011 [6].

The comparison of the omega 3 and omega 6 fatty acid contents of chia seeds with other seeds and grains has been given in fig. 2.




Figure 2: Omega 3 and Omega 6 fatty acid content of chia seeds and other seeds/grains – a comparison

3.4 Antioxidants

Chia seeds are high in phenolic compounds which have been scientifically proven to exhibit antioxidant related functions. Antioxidants and phenolic compounds have been found to have health promotive properties and also confer protection from degenerative diseases such as cardio vascular diseases, cancers, diabetes and diverticulosis [21]. Chia seeds and oil also contain several bioactive compounds namely quercetin, myricetin, kaempferol, chlorogenic acid and 3,4-dihydroxyphenylethanol-elenolic acid dialdehyde (DHPEA-EDA). Several in-vitro assays have confirmed that these polyphenols possess high antioxidant capacity and their presence is associated with lower levels of lipid autoxidation. Therefore, chia seeds can be used as a functional ingredient in the formulation of products for commercial applications. TABLE 5 gives the total phenol content of chia seeds as reported in different research studies [2, 5, 17, 22, 23,24].

TABLE 5: Mean total phenol content of Chia Seeds

Chia seeds of different origin	Jalisco, (Mexico)[5]	Sinaloa, (Mexico)[5]	Chia seeds (Chile) [15]
Total Phenols(gallic acid equivalent /g)	0.92 mg	0.88 mg	0.94 mg

3.5 Vitamins and Minerals

Chia has also been found to be a good source of several vitamins and minerals particularly niacin, zinc, calcium, phosphorus and magnesium. Niacin content of chia is higher than other cereals (corn, soybeans and rice), whereas the riboflavin and thiamin content is similar to that present in corn and rice. Chia contains six times more calcium, eleven times more phosphorus and four times more potassium than 100g of milk [16, 25].

TABLE 6: Average vitamins and minerals content of chia seeds

Vitamins (per 100g)		Minerals (per 100g)	
Ascorbic acid (mg)	1.6	Calcium (mg)	631
Thiamin (mg)	0.62	Iron (mg)	7.72
Riboflavin (mg)	0.17	Magnesium (mg)	335
Niacin(mg)	8.83	Phosphorus (mg)	860
Vitamin-A (IU)	54	Potassium (mg)	407
Vitamin-E (mg)	0.50	Potassium (mg)	407
		Sodium (mg)	16
		Zinc (mg)	4.58

Source: USDA National Nutrient Database for Standard Reference Release 28, 2011 [6].

IV. PHYSIOCHEMICAL PROPERTIES AND APPLICATIONS IN THE FOOD INDUSTRY

Moisture influences the structure and physical properties of chia seeds significantly. The physical properties of chia seeds have been studied by several researchers. The mean moisture content of white seeds is 7.2 per cent while that of black seeds is 6.6 per cent. The white seeds possess lower bulk density than black seeds, which could be attributed to their larger size. Bulk density signifies the storage and transport capacity of seeds. No significant difference has been found between the true density and porosity of both black and white seeds [1,16]. TABLE 7 gives the moisture density related physical properties of chia seeds.

TABLE 7: Moisture - dependent physical properties of black and white chia seeds

Physical property	Black seeds	White seeds
Bulk density (g cm ⁻³)	0.772 g cm ⁻³	0.667 g cm ⁻³
True density (g cm ⁻³)	1.009 g cm ⁻³	0.999 g cm ⁻³
Porosity (per cent)	28.2 per cent	33.1 per cent

Chia seeds have the unique property of forming a gelatinous mass when soaked in water. This is due to the presence of high amount of mucilages and gums. Studies indicate that chia seeds can absorb water up to 12 times their weight [16]. This property makes it useful in the food industry. Chia seeds could therefore also be friendly for the gut health because soluble fiber prolongs the gastro-intestinal transit time that aids in improved digestion. Dry chia seeds can be stored for longer period because of significantly higher concentration of natural antioxidants in them. Antioxidants prevent auto-oxidation of fatty acids present in chia-seed[26, 16].Chia seeds are exceptionally rich in mucilage. The mucilage of chia seeds contains about 71.22 per cent polysaccharides which is comparable to the polysaccharides in mucilage of flaxseeds (75per cent). But quite lower to that of mustard seeds (80 per cent) and xanthum gum (98 per cent) [27, 28, 29, 30]. Mucilage of chia seeds can be extracted by hydrating whole seeds at 20⁰C for 2 hours in the 1:40 seed and water ratio. After hydration a gelatinous solution gets formed, which can be collected and dried when kept at 50⁰ C for 10 hours in an air convection heat oven. Dried mucilages can be separated from the seeds over a 40 mesh screen via abrasion procedure. This extracted mucilage has been found to retain moisture up to 27 times of its weight on hydration, which shows its exceptionally high water holding capacity. It can also form highly viscous solution at lower

concentration (0.3 per cent) which is similar to xanthum gum. Solubility of chia mucilages has been found to be 100 per cent when examined at different concentrations i.e. 0.15, 0.25 and 0.5 per cent at temperatures (30, 60, 70 and 90^o C respectively) and also in different centrifugation (800 and 2000g) conditions [16]. Studies have indicated that, mucilages can replace oil and egg content up to 25g/100g in cake preparations; greater incorporation has been found to unfavorably alter the sensory properties of cake [31]. Another study suggests that chia mucilage can be used as an emulsifier and stabilizer in an ice cream after completing the texture, overrun and melting tests. Extracted mucilage of chia has been used to improve/maintain the quality of ice cream during storage [32].

Protein rich fractions of defatted chia flour produced by dry fractionation method have been found to exhibit good emulsifying properties as compared to foaming properties. Emulsion stability (95 per cent) of chia seed has been found to be higher at alkaline pH [33]. According to Coorey et al (2014), the water holding capacity (WHC: 266g of water/g of sample), oil holding capacity (OHC: 58.6g of water/g of sample) and emulsion stability (69.83 per cent) of extracted chia seed gel is found to be better than that of guar gum (WHC- 24.83g, OHC- 0.87g and emulsion stability-35.17 per cent) which is used for commercial purposes. Compared to the chia seeds gel, chia flour gel has been found to have good potential to be used as a thickening agent because of its functional properties and also because the former retains more amounts of omega-3 and omega-6 fatty acid after extraction. Freeze – thaw stability of chia seed gel makes it suitable as a commercially potential ingredient to be used in frozen products [34].

Attempts have been made to develop several products by incorporating chia seeds/ its flour or gel. In a study, whole chia flour was taken in four different proportions (5 per cent, 10 per cent, 12 per cent and 15 per cent) for preparing gluten-free, omega-3 and fibre rich chips. Chips with 5 per cent of chia seeds flour had highest consumer acceptability. No significant variations in the sensory characteristics (appearance, color, flavor and overall liking) were seen in the chips (5 per cent chia flour) when compared with commercially available ones [35]. Breads have also been prepared by incorporation of chia seeds. In the preparation of bread, when wheat flour is substituted with chia seeds (11.0g/100g) and chia flour (7.8g/100g) there is a resultant reduction in the saturated fatty acid (27 to 24 per cent) content. There is also an increase in the polyunsaturated-saturated fatty acid ratio in the bread with chia seed (288.1 per cent) and with chia flour (206.9 per cent). Breads prepared from chia seeds/chia flour are a good source of fiber and omega-3 fatty acids as compared to breads available in the market which are prepared from refined wheat flour/ whole wheat flour. It has been reported by Coelho et al (2015) that bread with chia flour has an acceptability index of 90 per cent. It was found that the consumer preference to buy such breads was 60 per cent in the “surely buy the product” category and 35 per cent in the category of “likely to buy the product”. Hence the preference for purchasing the breads with chia is fairly high [36]. Zettel et al (2014) prepared a gel of grounded chia seeds with 5g and 10g/g water respectively and concluded that dough with 2 per cent of chia gel tends to be softer in consistency. The quality of bread also improves in terms of storage, when mixed with 5g/g of water (optimal dosage). Breads with chia gel are found to be more nutritious and could be used as a functional food [37]. Inclusion of whole chia flour (0-20 per cent) and vital gluten (0-4 per cent) in bread for the development of functional product indicated that addition of 10 per cent whole chia flour and 2 per cent Vital Gluten would provide more amount of protein (19 per cent), ash

(11 per cent) and 26 per cent more lipids than standard loaf. However, the addition of whole chia flour in the dough reduces the protein network which results in a denser product [38].

Attempts have been made to incorporate chia in other bakery products as well. In a study, Inglett et al (2014) prepared four types of cookies – standard, chia - nutrim, chia - whole oat flour (WOF), chia - oat bran composite (OBC) to study the physical properties of chia –oat composites, dough and cookies. Addition of 20 per cent composite improved the Water-Holding Capacity (WHC) of cookies as compared to those prepared by the standard (non-chia) recipe. This study indicated that substituting wheat flour with Chia-Wheat oat flour and Chia-Oat bran composites will not possess any impact on texture of cookies. Fibre content and omega-3 enriched cookies proved to be very beneficial if used commercially [39]. Incorporation of the chia seeds in frankfurters was studied by Pintado et al (2016). Reformulated frankfurters with *Salvia hispanica* contained higher amount of protein, minerals, insoluble dietary fibers along with high levels of monounsaturated fatty acids and omega-3 fatty acids. [40].

The above discussed studies support the technological view point that, chia seeds possess many important functional properties (water-holding capacity, oil holding capacity, solubility, viscosity, emulsion stability and foaming stability) which prove its potential to be used as a thickening agent, gel forming agent, chelator, foam enhancer, emulsifying agent, clarifying agent, rehydrating agent and as suspension formers in the formulation of food products at both home and commercial level. All these properties make chia a promising functional food for the future.

V. EFFECTS ON HEALTH AND PREVENTION OF DISEASE

Several animal and human studies have been carried out to study the effects of consuming chia seeds on health. Some of the studies have been discussed below.

5.1 Effect of chia on animal health

Five thousand four hundred 1-day old male broiler chicks were fed on two different diets containing 10 per cent and 20 per cent of chia seeds for 49 days and at the end of the study, a remarkable fall in saturated fatty acid, polyunsaturated fatty acid: saturated fatty acid ratios and omega 6 – omega 3 ratios of white and dark meats were seen in comparison to meats of chicks fed on control diet [41]. Another trial was conducted by Coorey et al (2014) on 48 Isa Brown hens divided into 4 groups (six each). They were fed with 8 different diets [control diet, control with flax seeds (15g, 20g, 30g/100g of control), chia seeds (20g, 30g, 40g) and fish oil (1.5 ml/100g of control)] for 4 weeks. Researcher concluded that the omega-3 fatty acid content of egg yolk of hen that were fed with 30 per cent chia seeds was highest as compared to others without compromising the flavor of egg. Eggs with higher omega-3 fatty acid content may exhibit anti-hypercholesterolemic effects when consumed by humans [42].

Chicco and his colleagues (2009) examined the effect of alpha linolenic acid and fiber rich chia seed on dyslipidemia and insulin resistance in 72 Wistar rats, which were divided into 3 groups and assigned with three diets for three weeks in one experimental design. In another experimental design of the same study 96 rats which had been earlier fed with sucrose rich diet for 3 months were divided into two groups; one group continued with sucrose – rich diet while second group was provided with sucrose rich diet along with chia seeds.

It was noted that, chia seeds prevented the onset of dyslipidemia, peripheral insulin resistance and reduced the visceral adiposity in rats without producing changes in blood glucose levels [43]. Marineli et al (2015) conducted a trial on 36 Wistar rats (induced obesity) to examine the effect of chia seeds and oil on their plasma and liver oxidative status. Rats were divided into 6 groups and fed with two diets – one standard and another high fructose high fat diet (soybean oil, lard and fructose). Two treatments; one long treatment (12 weeks) and one small treatment (6 weeks), were followed for incorporation of chia seeds and oil in their diets. The feeding trial resulted in increased plasma catalase activity and increased plasma thiol (GSH) as compared to high fructose high fat diet. No significant effect was observed on the liver lipid peroxidation [44]. In yet another study, 48 Wistar rats were fed with an innovatively developed omega-3 enriched chocolate milk and its effect was examined on the muscle damage and blood lipid profile. Reduction in the muscle damage with lower levels of total cholesterol and triacylglycerols were seen which indicate that intake of omega-3 enriched beverage alters the post exercise soreness of muscle. No side-effects were noted on lungs, spleen, liver, heart and kidney after ingesting the beverage for 15 days [45].

5.2 Effect of Chia on Human health

Very few studies have been done on human volunteers to assess the impact of consuming chia seeds on health. A randomized, placebo-controlled clinical trial on 76 adults found no significant reduction in body weight, lipid profile, inflammatory markers and blood sugar levels even after ingesting 50g chia seeds per day for 12 weeks. Elevation in plasma alpha linolenic acid (24.4 per cent) was noted as compared to the control group [46]. Effect of ingesting 25g milled chia seeds for seven weeks was examined on 10 postmenopausal healthy women. The ingestion of chia seeds resulted in significant increase in the eicosapentaenoic acid (EPA) and alpha linolenic acid (ALA) levels in plasma by 30 per cent and 138 per cent respectively. However, no significant change was seen on body mass index [47]. Higher intake of omega-3 fatty acids in diet through both marine (EPA and DHA) and plant sources (ALA) are strongly associated with decreasing the risk of death due to cardiovascular diseases which is a major cause of mortality in the world nowadays [19].

A randomized, placebo-controlled clinical trial was conducted by Vuksan et al (2007) on twenty type-2 controlled diabetics (with medication) who were provided 37g chia seeds daily (added in white bread). The results indicated that high fiber content of chia helped in controlling hyperglycemia and reducing systolic blood pressure. ALA and EPA levels were found to increase with consumption of chia. Anticoagulant and anti-inflammatory effect of chia seeds may help in preventing strokes and heart attacks in type-II diabetic patients [48]. Few studies indicate that chia seeds exhibit positive effects on health such as reduction of post-prandial blood glucose levels and systolic blood pressure [49, 18, 50]. In a double-blind placebo controlled trial sixty seven adults (divided into two groups) with excess body weight and metabolic syndrome were given 4g chia seeds mixed with oats, palm and soy powder diluted in 250 ml of water two times per day for two months along with some caloric modifications in the daily diet. The study results indicated a significant reduction in the body weight, waist circumference and basal metabolic rate of experimental group in comparison to placebo group. Significant reduction in insulin resistance, triglycerides levels and C-reactive protein were noted. However, no significant change was seen in the total cholesterol, blood sugar and total plasma insulin levels [21].

VI. SIDE EFFECTS OF CHIA SEEDS

European Parliament and Council of Europe approved chia seeds as a Novel Food in 2009. Till date no study has revealed any adverse toxic, allergic or anti-nutritional effects after ingesting whole or ground chia seeds. The food safety and standards act (2006) and its regulations (2011) do not provide any information regarding the prohibition of chia seeds. Chia seeds may thus be considered safe for human consumption and has a promising future as functional food [51, 52, 53]. However, more clinical trials on animals and humans need to be done to ascertain the safety aspects of chia seeds.

VII. CONCLUSION

Chia seeds are not new to mankind. They have been used even in the pre-colombian time by Aztecs as foodstuff and in religious ceremonies. Chia seeds are an excellent source of dietary fiber (insoluble and soluble), omega-3 fatty acids, proteins and bioactive compounds or phytochemicals. Chia possesses many important physiochemical and functional properties which makes it more suitable in the food industry. Chia acts as a good thickener, gel former, chelator, foam enhancer, emulsifier, suspension formers, clarifying agent and as a rehydrating agent. Therefore it can be used commercially for the development of new products enriched with omega-3, protein, soluble/insoluble fiber and phenolic compounds. Chia seeds may help in the prevention, treatment and management of several non-communicable diseases, improving immunity and perhaps modifying the blood clotting mechanism. Chia also helps in improving the post-prandial blood glucose levels in blood by slowing down the digestion of carbohydrates. It can be incorporated in frozen products, bakery, beverages, sweets, baby foods, pasta, sausages etc. Researches done in *vivo* and *vitro* have supported the fact that that it is safe for human consumption and also exhibits wide range of health benefits. There is scope for research on chia seeds with respect to the food industry and nutraceuticals. Chia can thus be considered as a functional food which could help in improving the health of the masses.

REFERENCES

- [1] V. Y. Ixtainaa, S. M. Nolasco, M. C. Tomas, Physical properties of chia (*Salvia hispanica* L.) seeds, *Journal of Industrial crops and products* 28 (3), 2008, 286–293.
- [2] M.I. Capitani, V. Spotorno, S.M. Nolasco, M.C. Tomás, Physicochemical and functional characterization of by-products from chia (*Salvia hispanica* L.) seeds of Argentina, *LWT - Food Science and Technology*, 45, 2012, 94-102.
- [3] R. Ayerza, W. Coates, Chia. Rediscovering a Forgotten Crop of the Aztecs, (The University of Arizona Press, 1st edition, Tucson, Arizona, p. 197, 2005).
- [4] W. Coates. Whole and ground chia (*Salvia hispanica* L.) seeds, chia oil - effects on plasma lipids and fatty acids. *Nuts and Seeds in Health and Disease Prevention*, 2011, DOI: 10.1016/B978-0-12-375688-6.10037-4.
- [5] E. Reyes-Caudillo, A. Tecante, M.A. Valdivia-Lo'pez, Dietary fiber content and antioxidant activity of phenolic compounds present in Mexican chia (*Salvia hispanica* L.) seeds, *Journal of Food Chemistry*, 107, 2008, 656–663.

- [6] USDA National Nutrient Database for Standard Reference Release 28 (2011). Basic report 12006, seeds, Chia seeds, dried. Report date: January 11, 2016
- [7] C. Gopalan, B.V. Rama Sastri and S.C. Balasubramanian, Nutritive value of Indian foods(National Institute of Nutrition, Indian Council of Medical Research, Hyderabad, India, 2007).
- [8] R. Ayerza, and W. Coates, Composition of chia (*salvia hispanica*) grown in six tropical and subtropical ecosystems of South America, *Journal of Tropical Science*, 44(3), 2004, 131-135.
- [9] R. Ayerza and W. Coates, Protein content, oil content and fatty acid profiles as potential criteria to determine the origin of commercially grown chia (*Salvia hispanica* L.), *Journal of Industrial Crops and Products*, 34, 2011, 1366– 1371.
- [10] R. Ayerza and W. Coates, Chia seeds: New source of omega-3 fatty acids, natural antioxidants and dietary fiber, (South-East Center for Natural Products Research and Commercialization, Office of Arid lands studies, Tucson, Arizona, USA, 2001)
- [11] J. V. Ovando, G. R. Rubio, L. C. Guerrero, and D. B. Ancona, Physicochemical properties of fibrous fraction from chia (*Salvia hispanica* L.), *LWT- Journal of Food Science and Technology*, 42, 2009, 168-173.
- [12] S. Suri, Effect of Dietary fiber (Guar-Gum) supplementation on hypercholesterolemic patients with or without hyperglycemia, M.Sc. dissertation, Institute of Home Economics, University of Delhi, 1996.
- [13] J. W. Anderson, P. Baird, R. H. Davis, S. Ferreri, M. Knudtson, A. Koraym, Health benefits of dietary fiber, *Nutrition Reviews*, 67(4), 2009, 188-205.
- [14] S. Suri, S.J. Passi and S. C. Manchanda, Effect of prebiotic – guar gum supplementation among dyslipidemic patients with or without hyperglycemia, *International Journal of Advanced Technology in Engineering and Science*, 3(11), 2015.
- [15] W. Coates and R. Ayerza, Chia (*Salvia hispanica* L.) seeds as an omega-3 fatty acid source for feeding pigs: effects on fatty acid composition and fat stability of the meat and internal fat, growth performance and meat sensory characteristics, *Journal of Animal Science*, 87, 2009, 3798-3804, doi: 10.2527/jas.2009-1987.
- [16] L.A. Munoz, A. Cobos, O. Diaz, and J. M. Aguilera, Chia seeds: microstructure, mucilage extraction and hydration, *Journal of Food Engineering*, 108, 2012, 216-224.
- [17] R. Da Silva Marineli , E. A. Moraes , S. A. Lenquiste , A. T. Godoy , M. N. Eberlin , M. R. Maróstica Jr., Chemical characterization and antioxidant potential of chilean chia seeds and oil (*Salvia hispanica* L.) , *LWT - Food Science and Technology*, 59, 2014, 1304-1310. <http://dx.doi.org/10.1016/j.lwt.2014.04.014>
- [18] L.T. Toscano, C.S. Oliveira Da Silva, L.T. Toscano and A.E. Monteiro De Almeida and A.D. Cruz santos and S.S. Alexandre, Chia flour reduces blood pressure in hypertensive subjects, *Journal of Plant Foods and Human Nutrition*, 69, 2014, 392-398.
- [19] A. S. Koh,A. Pan,R. Wang,A. O. Odegaard,M. A. Pereira,J. M. Yuan, and W. P. Koh, The association between dietary omega-3 fatty acids and cardiovascular death: the Singapore Chinese health study, *European Journal of Preventive Cardiology*, 22(3), 2015, 364–372. Doi: 10.1177/2047487313517576.

- [20] I. R. Ayerza, Chia as an Omega-3 fatty acid source for animal and human consumption, Office of Arid Lands Studies, The University of Arizona. <https://azchia.com/research/Omega3source.htm> (report date 1/9/2016), cited on 18/03/2016.
- [21] M. Guevara- Cruz, A. R. Tovar, C. A. Aguilar- Salinas, I. Medina – Vera, L. Gil-Zenteno, I. Hernandez-Viveros, P. Lopez-Romero, G. Ordaz-Nava, S. Canizales-Quinteros, L. E. Guillen Pineda, and N. Torres, A dietary pattern including nopal, chia seeds, soy protein and oat reduces serum triglycerides and glucose intolerance in patients with metabolic syndrome, *The Journal of Nutrition and Disease*, 142, 2012, 64-69, doi:10.3945/jn.111.147447.
- [22] L.M. Alvarez- Chavez, M.A. Valdivia-Lopez, M. L. Aburto -Juarez, and A. Tecante, Chemical characterization of the lipid fraction of Mexican chia seed (*Salvia hispanica* L.). *International Journal of Food Properties*, 11, 2008, 687-697.
- [23] V. Y. Ixtaina, M. L. Martinez, V. Spotorno, C. M. Mateo, D. M. Maestri, B. W.K. Diehl, Characterization of chia seed oils obtained by pressing and solvent extraction. *Journal of Food Composition and Analysis*, 24, 2011, 166-174.
- [24] O. Martinez – Cruz and O. Parades-Lopez, Phytochemical profile and Nutraceutical potential of chia seeds (*Salvia hispanica* L.) by ultra high performance liquid chromatography, *Journal of Chromatography A*, 1346, 2014, 43-48.
- [25] L. H. Munoz, Mucilage from chia seeds (*salvia hispanica* L.): microstructure, physico-chemical characterization and applications in food industry, doctoral diss., Pontificia Universidad Catolica De Chile Escuela De Ingenieria, Santiago de Chile, 2012.
- [26] V.Y. Ixtaina, S.M. Nolasco, M.C. tomas, Oxidative stability of chia (*salvia hispanica* L.) seed oil: effect of antioxidants and storage conditions, *Journal of American Oil Chemists Society*, 89, 2012, 1077-1090.
- [27] J. Warrand, P. Michand, P. Picton, L. Muller, G. Curtosis, B. Ralainirina, R. Curtosis, Structural investigation of the neutral polysaccharide of *Linum usitatissimum* L. seeds mucilage, *International Journal of Biological Macromolecules*, 35(3-4), 2005, 121-125.
- [28] W. Cui, N.A.M. Eskin, C.G. Biliaderies, Chemical and physical properties of yellow mustard (*Sinapsis alba* L.) mucilage, *Journal of Food Chemistry*, 46(2), 1993, 169- 176.
- [29] M. Ahmed, R. Hamed, M. Ali, A. Hassan, E. Babiker, Proximate composition, antinutritional factors and protein fractions of guar gum seeds as influenced by processing treatments, *Pakistan Journal of Nutrition*, 5(5), 2006, 340-345.
- [30] S. Suri and A. Malhotra, *Food science, nutrition and safety* (Dorling Kindersley, India, Pvt. Ltd, 2014).
- [31] R. Borneo, A. Aguirre, A. E. León, Chia (*Salvia hispanica* L) gel can be used as egg or oil replacer in cake formulations, *Journal of the American Dietetic Association*, 110, 2010, 946-949.
- [32] B.E. Campos, T. Dias Ruivo, M.R. da Silva Scapim, G.S. Madrona, R.de C. Bergamasco, Optimization of the mucilage extraction process from chia seeds and application in ice cream as a stabilizer and emulsifier, *LWT - Food Science and Technology*, 2015, doi: 10.1016/j.lwt.2015.09.021.
- [33] A. Vázquez-Ovando, D. Betancur-Ancona and L. Chel-Guerrero, Physicochemical and functional properties of a protein-rich fraction produced by dry fractionation of chia seeds (*Salvia hispanica* L.), *CyTA - Journal of Food*, 11:1, 2013, 75-80, <http://dx.doi.org/10.1080/19476337.2012.692123>.

- [34] R. Coorey, A. Tjoe and V. Jayasena, Gelling properties of chia seed and flour, *Journal of Food Science*, 79(5), 2014.
- [35] R. Coorey, A. Grant and V. Jayasena, Effects of chia flour incorporation on the nutritive quality and consumer acceptance of chips, *Journal of Food Research*, 1(4), 2012, <http://dx.doi.org/10.5539/jfr.v1n4p85>.
- [36] Michele Silveira Coelho, Myriam de las Mercedes Salas-Mellado. Effects of substituting chia (*Salvia hispanica* L.) flour or seeds for wheat flour on the quality of the bread, *LWT - Food Science and Technology* 60 (2015) 729-736
- [37] V. Zettel, A. Kramer, F. Hecker, B. Hitzmann, Influence of gel from ground chia (*Salvia hispanica* L.) for wheat bread production. *Europe Food Research Technology*, 2014, DOI: 10.1007/s00217-014-2368-8
- [38] Patricia Luna Pizarro, Eveline Lopes Almeida, Alessandra Silva Coelho, Norma Cristina Samman, Miriam DupasHubinger, Yoon Kil Chang (2014) Functional bread with n-3 alpha linolenic acid from whole chia (*Salvia hispanica* L.) flour. *Journal of Food Science and Technology*. DOI: 10.1007/s13197-014-1477-5.
- [39] George E Inglett, Diejun Chen and Sean Liu. Physical properties of sugar cookies containing chia–oat composites, *J Sci Food Agric* 2014; 94: 3226–3233.
- [40] T. Pintado, A.M. Herrero, F. Jiménez-Colmenero, C. Ruiz-Capillas. Strategies for incorporation of chia (*Salvia hispanica* L.) in frankfurters as a health-promoting ingredient, *Meat Science* 114 (2016) 75–84.
- [41] R. Ayerza, W. Coates and M. Lauria, Chia seed (*Salvia hispanica* L.) as an omega-3 fatty acid source for broilers: influence on fatty acid composition, cholesterol and fat content of white and dark meats, growth performance and sensory characteristics, *Journal of Poultry Science*, 81, 2002, 826-837.
- [42] R. Coorey, A. Novinda, H. Williams and V. Jayasena, Omega-3 fatty acid profile of eggs from laying hens fed diets supplemented with chia, fish oil and flaxseed, *Journal of Food Science*, 2014, doi:10.1111/1750-3841.12735.
- [43] A. G. Chicco, M.E. D. Alessandro, G. J. Hein, M. E. Oliva and Y. B. Lombardo, Dietary chia seed (*Salvia hispanica* L.) rich in alpha linolenic acid improves adiposity and normalizes hypertriglycerolaemia and insulin resistance in dyslipaemic rats, *British Journal of Nutrition*, 101, 2009, 41-50.
- [44] R. D. S. Marinelli, S.A. Lenquiste, E. A. Moraes, M. R. Marostica Jr., Antioxidant potential of dietary chia seed (*Salvia hispanica* L.) and oil in diet induced obese rats, *International Journal of Food Research*, 2015, doi: 10.1016/j.foodres.2015.07.039.
- [45] P. N. Morato , J. B. Rodrigues , C. S. Moura , F. G. Drummond e Silva , E. A. Esmerino , A. G. Cruz , H. M. A. Bolini , J. Amaya-Farfan , P. C. B. Lollo, Omega-3 enriched chocolate milk: A functional drink to improve health during exhaustive exercise, *Journal of Functional Foods*, 14, 2015, 676–683.
- [46] D. C. Nieman, E.J. Cayea, M. D. Austin, D.A. Henson, Mc. S. R. Anulty, F. Jin, Chia seed does not promote weight loss or alter disease risk factors in overweight adults. *Journal of Nutrition Research*, 29, 2009, 414-418.
- [47] F. Jin, D. C. Nieman, E. Sha, G. Xie, Y. Qiu and W. Jia, Supplementation of milled chia seeds increases plasma ALA and EPA in postmenopausal women, *Journal of Plant Food and Human Nutrition*, 67, 2012, 105-110.

- [48] V. Vuksan, D. Whitman, J. Sievenpiper, A. Jenkins, A. Rogovik, R. Bazinet, E. Vidgen and A. Hanna, Supplementation of conventional therapy with the novel grain Salba(*Salvia hispanica* L.) improves major and emerging cardiovascular risk factors in type-2 diabetes, *Journal of Diabetes Care*, 30(11) 2007, 2804-2810.
- [49] V. Vuksan, A.L., Jenkins, A. G. Dias, A.S. Lee, E. Jovanovski, A. L. Rogovik and A. Hanna, Reduction in postprandial glucose excursion and prolongation of satiety: possible explanation of the long – term effects of whole grain Salba(*Salvia hispanica* L.), *European Journal of Clinical Nutrition*, 64(4),2010, 436-438.
- [50] H. Ho, A.S. Lee, E. Jovanovski, A. L. Jenkins, R. Desouza, V. Vuksan, Effect of whole, ground salba seeds (*Salvia hispanica* L.) on post-prandial glycemia in healthy volunteers: a randomized controlled, dose –response trial, *European Journal of Clinical Nutrition*, 67(7), 2013, 786-788.
- [51] Commission, E.U., 2009. Authorizing the placing on the market of chia seeds (*Salvia hispanica*) as novel food ingredient under Regulation (EC) No. 258/97 of European Parliament of the Council. *Official Journal of the European Union*, C, 7645.
- [52] EFSA “opinion of the scientific panel on dietetic products, nutrition and allergies on a request from the European commission related to the safety of chia (*Salvia hispanica* L.), Seeds and ground whole chia as a novel food ingredient intended for use in bread,” *EFSA Journal*, 278, 2005, 1-12.
- [53] EFSA, “scientific opinion of the panel on dietetic products, nutrition and allergies on a request from the European commission on the safety of ‘chia seeds (*Salvia hispanica* L.)and ground whole chia seeds’ as a food ingredient”, *The EFSA Journal*, 996, 2009, 1-2.