

## ON H-ADHERENCE AND MINIMAL $S_{2\frac{1}{2}}$ SPACES

Anakh Singh

Assistant Professor, CGC, Landran, Mohali, (India)

### ABSTRACT

In this paper we study the H-adherence based in terms of hyperclosure, of a filterbase obtain characterizations of some of the spaces in terms of H-adherence. Also variant of Minimal-P spaces and its relationships with other axioms is investigated.

**2010 Mathematics Subject Classification:** 54A10, 54A20, 54D10, 54D25, 54D30.

**Key words:** H-adherence, Urysohn closed,  $U(i)$ , Minimal Urysohn, Minimal  $S_{2\frac{1}{2}}$ .

### I INTRODUCTION

The adherence of a filterbase has been used to characterize some of the well known topological spaces such as compact spaces, H-closed spaces and various minimal topological spaces. Various forms of adherence of filterbase have been discussed in literature and these have found much importance place in topology. The minimal and maximal topologies with respect to a certain topological property in the lattice of topologies  $LT(X)$  defined on a set  $X$  partially ordered by inclusion have been of keen interest of topologists. In [1] Berri et.al, have given a survey on the considerable amount of work done in the field of minimal topological spaces. In [2] Herrington, introduced u-adherence of a filterbase and characterized Urysohn closed and Minimal Urysohn spaces in terms of u-adherence.

In this paper we study a variant of adherence based on hyperclosure of a set, namely H-adherence of a filterbase and characterize some spaces in terms of H-adherence. Also we investigate the class of minimal topological spaces with respect to the separation axiom  $S_{2\frac{1}{2}}$  and the role of H-adherence in characterization of these spaces.

**Notations:** Throughout the paper, by a space  $X$  we mean a topological space,  $N_{\mathfrak{F}}(x)$  is the filterbase of  $\mathfrak{F}$ -neighborhoods of some  $x \in X$ ,  $cl_{\mathfrak{F}}(A)$  denotes the closure of the subset  $A \subset X$ ,  $X \setminus A$  the complement of  $A$  in  $X$  and  $\mathbf{Z}$  denotes the set of integers.

**Definition 1.1:** (1) A filter base  $B$  on a space  $X$  is said to be an open filter if each member of  $B$  is an open set.

(2) A point  $x$  is called a *adherent point* or *accumulation point* or *cluster point* of a filterbase  $B$  denoted as  $x \in \text{adh } B$ , if for each open sets  $U$  containing  $x$  and each  $F$  in  $B$ ,  $F \cap U \neq \emptyset$ .

(3) A filterbase  $B$   $\theta$ -converges to  $x$  if for each open set  $U$  containing  $x$  there exists an  $F$  in  $B$  such that  $cl(F) \subset U$ .

(4) A point  $x$  is called a  $\theta$ -adherent point of a filterbase  $B$  denoted as  $x \in \text{adh}_\theta B$ , if for each open sets  $U$  containing  $x$  and each  $F$  in  $B$ ,  $F \cap cl(U) \neq \emptyset$ .

**Definition 1.2 [3]:** If  $X$  is a space,  $A \subset X$ , and  $x \in X$ , then

(1)  $\ker(A) = \bigcap \{ U : U \in \mathfrak{F} \text{ and } A \subset U \}$ .

(2)  $\langle x \rangle = cl\{x\} \cap \ker\{x\}$ .

(3)  $cl_H(A) = \{x \in X : \text{For each open set } U \text{ containing } x \text{ and each open set } V \text{ containing } A, cl(U) \cap cl(V) \neq \emptyset\}$

(4) A is *hyperclosed* if  $A = \text{cl}_H(A)$ .

**Definition 1.3 [4]:**(1) An open filterbase B is a *Urysohn filterbase* if and only if for each x not in adherence of B, there is an open set U containing x and V in B such that,  $\text{cl}(U) \cap \text{cl}(V) = \emptyset$ .

(2) A pair of open sets G and H containing a point x is called an ordered pair denoted by

(G, H) if,  $x \in G \subset \text{cl}(G) \subset H$ .

(3) For any subset  $A \subset X$ ,  $\text{cl}_u(A) = \{x \in X: \text{For each ordered pair of open sets (G, H) containing x, } A \cap \text{cl}(H) \neq \emptyset\}$ .

(4) A filterbase B *u-converges* to x if for each ordered pair of open sets (G, H) containing x there exists an F in B such that  $F \subset \text{cl}(H)$ .

(5) A point x is called a *u-adherent point* or *u-accumulation point* of a filterbase B denoted as  $x \in \text{adh}_u B$ , if for each ordered pair of open sets (G, H) containing x and each F in B,  $F \cap \text{cl}(H) \neq \emptyset$ .

(6) An Urysohn space X is *Urysohn-closed* provided X is a closed set in every Urysohn space in which it can be embedded.

(7) A space  $(X, \mathfrak{T})$  is said to be *Minimal Urysohn* if  $\mathfrak{T}$  is Urysohn and there exists no Urysohn topology on X strictly weaker than  $\mathfrak{T}$ .

**Definition 1.4 [5]:** A space X is said to be,

(1) *Urysohn ( $T_{2\frac{1}{2}}$ ) space* if for every pair of points x and y in X there exist neighborhoods U of x and V of y such that  $\text{cl}(U) \cap \text{cl}(V) = \emptyset$ .

(2)  *$S_{2\frac{1}{2}}$  space* if for every pair of points x and y in X, whenever  $\text{cl}\{x\} \neq \text{cl}\{y\}$  then there exist neighborhoods U of x and V of y such that  $\text{cl}(U) \cap \text{cl}(V) = \emptyset$ .

(3)  *$S_2$  space* if for every pair of points x and y in X, whenever  $\text{cl}\{x\} \neq \text{cl}\{y\}$  then there exist disjoint neighborhoods containing them.

(4)  *$S_1$  space* if for every pair of points x and y, whenever x has a neighborhood not containing y, then y has a neighborhood not containing x.

**Definition 1.5:** A space X is said to be,

(1)  *$\theta$  - point paracompact [6]* if for each open covering  $\mathcal{U}$  of X and each  $x \in X$  there exists an open refinement  $\mathcal{V}$  of  $\mathcal{U}$  and a  $\theta$ -open set U containing x which intersects with only finite members of  $\mathcal{V}$ .

(2) H(i) [2] if every open filter on X has non void adherence.

(3) U(i) [2] if every Urysohn filterbase on X has non void adherence.

The following results will be used in the next section.

**Lemma 1.6:** A filter F  $\theta$ -adheres to a point x if and only if there exists a finer filter G which  $\theta$ -converges to the point x.

**Lemma 1.7 [7]:** A space  $X$  is  $S_1$  if and only if one of the following conditions is satisfied:

- (a) If  $U$  is open in  $X$  and  $x \in U$  then  $\text{cl}\{x\} \subset U$ .
- (b) If  $x, y \in X$ , then  $\text{cl}\{x\} = \text{cl}\{y\}$  or  $\text{cl}\{x\} \cap \text{cl}\{y\} = \emptyset$ .

**Lemma 1.8:** For a space  $X$  and  $x, y \in X$ , following statements are equivalent:

- (a)  $X$  is  $S_{2\frac{1}{2}}$ .
- (b) Either,  $\langle x \rangle = \langle y \rangle$  or  $\text{cl}_H\{x\} \cap \text{cl}_H\{y\} = \emptyset$ .

**Lemma 1.9:** For a space  $X$  and  $x, y \in X$ ,  $\langle x \rangle = \langle y \rangle$  if and only if  $\text{cl}\{x\} = \text{cl}\{y\}$ .

**Lemma 1.10:** Let  $X$  be Urysohn space. Then  $X$  is Urysohn closed if and only if every open filterbase has  $u$ -adherent point.

**Lemma 1.11:**  $X$  is  $U(i)$  if and only if every open filterbase has  $u$ -adherent point.

## II H-ADHERNCE

First we give the relationships between  $H$ -adherence and some forms of adherence of filterbases known in literature. For this purpose, we prove that  $u$ -closure and hyper-closure of a set coincide.

**Lemma 2.1:** For a space  $X$  and a subset  $A$  of  $X$ ,  $\text{cl}_u A = \text{cl}_H A$ .

**Proof:** Let  $x \in X$ ,  $x \in \text{cl}_H A \setminus \text{cl}_u A$ . Then there exists an ordered pair of open sets  $(G, H)$  containing  $x$  such that,  $A \cap \text{cl}(H) = \emptyset$ . Then,  $V = X \setminus \text{cl}(H)$  is an open set containing  $A$ , such that  $\text{cl}(V) \cap H = \emptyset$ . As  $(G, H)$  is an ordered pair containing  $x$ , we have  $\text{cl}(G) \cap \text{cl}(V) = \emptyset$  and this implies that  $x \notin \text{cl}_u A$  which is a contradiction.

Conversely, let  $p \in \text{cl}_u A \setminus S$ . Then there exists an open set  $U$  containing  $p$  and an open set  $V$  containing  $A$  such that,  $\text{cl}(U) \cap \text{cl}(V) = \emptyset$ . Thus  $(U, X \setminus \text{cl}(V))$  is an ordered pair of open sets containing  $p$  such that  $V \cap (X \setminus \text{cl}(V)) = \emptyset$ . Therefore,  $A \cap (\text{cl}(X) \setminus \text{cl}(V)) = \emptyset$  which implies  $x \notin \text{cl}_u A$  which is a contradiction.

**Definition 2.2:** A point  $x$  is called a  $H$ -adherent point of a filterbase  $B$  denoted as  $x \in \text{adh}_H B$ , if for each open set  $U_x$  containing  $x$  and each open set  $V_F$  containing  $F$ ,  $\text{cl}(U_x) \cap \text{cl}(V_F) \neq \emptyset$  for all  $F$  in  $B$ .

**Theorem 2.3:** For a space  $X$  and filterbase  $B$  of  $X$  the following hold:

- (a)  $\text{adh}_H B = \bigcap \{\text{cl}_H(F) \mid F \in B\}$ .
- (b)  $\text{adh}_u B = \text{adh}_H B$ .

**Proof:** Part (a) is straightforward. Part (b) follows from part (a) and Lemma 2.1, since  $\text{adh}_u B = \bigcap \{\text{cl}_u(F) \mid F \in B\}$ .

**Theorem 2.4:** For a space  $X$  and a subset  $A$  of  $X$ , the following are equivalent:

- (a)  $x \in \text{cl}_H(A)$ .
- (b) There exists a filter  $F$  such that  $A \in F$  and  $x \in \text{adh}_H F$ .

(c) There exists a filter  $F$  which  $\theta$ -converges to both  $x$  and  $A$ .

(d) There exists a filter  $F$  which  $\theta$ -adheres to both  $x$  and  $A$ .

**Proof:** (a)  $\Rightarrow$  (b) Let  $x \in \text{cl}_H(A)$  and  $B = \{A\}$  be a filterbase and  $F(B) = \{F \mid A \subset F\}$  be the filter generated by  $B$ . Now  $x \in \text{adh}_H B$  which implies  $x \in \text{adh}_H F(B)$ .

(b)  $\Rightarrow$  (c) Let  $F$  be filter such that  $A \in F$  and  $x \in \text{adh}_H F$ . Let  $P$  be a filterbase where,  $P = \{\text{cl}(U_F) \mid U_F \text{ are open sets containing } F \text{ for all } F \in F\}$ . Now as  $x \in \text{adh}_H F$  the  $G(P)$  the filter generated from the filterbase  $P$ ,  $\theta$ -adheres to  $x$ . By Lemma 1.6 there exists a finer filter  $R$  which  $\theta$ -converges to  $x$ . As  $A \in F$ ,  $\{\text{cl}(U_A) \mid U_A \text{ are open sets containing } A\} \subset P \subset G \subset R$ . Hence the filter  $R$   $\theta$ -converges to  $A$  also.

(c)  $\Rightarrow$  (d) is obvious

(d)  $\Rightarrow$  (c) follows from Lemma 1.6

(c)  $\Rightarrow$  (a) Let  $F$  be a filter which  $\theta$ -converges to both  $x$  and  $A$ . Then for every open set  $U$  containing  $x$  and every open set  $V$  containing  $A$  there exists  $F_1$  and  $F_2$  in  $F$  such that  $F_1 \subset \text{cl}(U)$  and  $F_2 \subset \text{cl}(V)$ . Since  $F$  is a filter  $F_1 \cap F_2 \in F$  and  $F_1 \cap F_2 \subset \text{cl}(V) \cap \text{cl}(U)$  and hence  $\text{cl}(V) \cap \text{cl}(U) \neq \emptyset$ . Thus  $x \in \text{cl}_H(A)$ .

**Corollary 2.5:** A set  $A$  of  $X$  is hyperclosed if and only if  $A$  contains all the  $H$ -adherent points of every filter containing  $A$  as a member.

As it is known that every  $H(i)$  space is  $U(i)$ , we give a condition in terms of  $H$ -adherence of open filters for which the converse holds.

**Lemma 2.6:** If  $X$  is  $\theta$ -point paracompact then every open filter with  $H$ -adherent point has an adherent point.

**Proof:** Let  $X$  be a  $\theta$ -point paracompact space and  $F$  be an open filter with an  $H$ -adherent point  $x$  and no adherent point. Then,  $\mathcal{U} = \{X \setminus \text{cl}(F) : F \in F\}$  is a directed open cover of  $X$  and so there exists an open refinement  $\mathcal{V}$  of  $\mathcal{U}$  and a  $\theta$ -open set  $K$  containing  $x$  which intersects with only finite members of  $\mathcal{V}$ . Let  $P = \cup\{U \in \mathcal{V} : U \cap K = \emptyset\}$ . Hence,  $P \cap K = \emptyset$ . Now,  $X \setminus P \subseteq \cup\{U \in \mathcal{V} : U \cap K \neq \emptyset\}$  and as  $\mathcal{V}$  is an open refinement of  $\mathcal{U}$  there exists finitely many  $O_i$  in  $\mathcal{U}$  such that  $\cup\{U \in \mathcal{V} : U \cap K \neq \emptyset\} \subseteq \bigcup_{i=1}^n \{O_i : O_i \in \mathcal{U}\}$ . Since  $\mathcal{U}$  is an open directed cover there exists some  $F \in F$  such that  $\bigcup_{i=1}^n \{O_i : O_i \in \mathcal{U}\} \subseteq X \setminus \text{cl}(F)$ . Thus  $X \setminus P \subseteq X \setminus \text{cl}(F)$  which implies  $\text{cl}(F) \subseteq P$ . Therefore we have,  $\text{cl}(F) \cap K = \emptyset$  and as  $K$  is a  $\theta$ -open set, by definition there exists an open set  $Q$  containing  $x$  such that  $\text{cl}(Q) \subset K$ . So,  $\text{cl}(Q) \cap \text{cl}(F) = \emptyset$  and as  $F$  being a member of an open filter is an open set we have  $x \notin \text{cl}_H(F)$  which implies that  $x$  is not an  $H$ -adherent point of  $X$ , a contradiction.

**Theorem 2.7:** A  $\theta$ -point paracompact space  $X$  is  $H(i)$  if and only if it is  $U(i)$ .

**Proof:** As  $H(i)$  space always implies  $U(i)$ , the result follows from Lemma 2.6 and Lemma 1.11 above.

### III MINIMAL $S_{2\frac{1}{2}}$ SPACES

**Definition 3.1[1] :** Given a topological property  $P$  in the lattice  $LT(X)$  on a set  $X$  a topology is said to be *minimal* if every weaker topology in  $LT(X)$  does not possess that property.

In this context of Minimal spaces we define Minimal  $S_{2\frac{1}{2}}$  space and characterize them in terms of  $H$ -adherence,

**Definition 3.2:** A topological space  $(X, \mathfrak{T})$  is said to *Minimal  $S_{2\frac{1}{2}}$  space* if  $\mathfrak{T}$  is the minimal element of the lattice  $LT(X)$  which is  $S_{2\frac{1}{2}}$ .

**Remark 3.3 [5]:** The separation axioms like  $S_{2\frac{1}{2}}$  which are non  $T_1$  are vacuously satisfied by the indiscrete topology and in such cases of separation axioms the study of minimal topological in the lattice  $LT(X)$  becomes trivial. So it becomes necessary to restrict the interval of topologies in the lattice  $LT(X)$  to avoid such circumstances.

So to each  $\rho \in LT(X)$  we associate the interval as in [2],  $L_\rho = \{ \mathfrak{T} \in LT(X) : at(\rho) \leq \mathfrak{T} \leq \overline{\rho} \}$  where  $at(\rho)$  denotes the topology on  $X$  generated by the sets  $\{X \setminus cl_\rho(P) : P \text{ is a finite subset of } X\}$  and  $\overline{\rho}$  denotes the closure of  $\rho$  in the power set  $2^X$ .

**Remark 3.4 [5]:** (1) If  $\rho \in LT(X)$  then  $cl_{at(\rho)}\{x\} = cl_\rho\{x\} = cl_{\overline{\rho}}\{x\}$  for all  $x \in X$ .

(2) If  $\mathfrak{T}, \rho \in Lt(X)$  then  $\mathfrak{T} \in L_\rho$  if and only if  $cl_\rho\{x\} = cl_{\mathfrak{T}}\{x\}$  for all  $x \in X$ .

(3) If  $\rho$  is  $S_1$  and if  $X$  can be written as finite disjoint point closures then for each point  $x \in X$ ,  $cl\{x\} \in at(\rho)$  which leads to the consequence  $at(\rho) = \rho = \overline{\rho}$  and thus  $L_\rho = \{\rho\}$ . To avoid this condition of triviality, in the further Section we assume that  $\rho \in LT(X)$  is an  $S_1$  topology such that  $X$  can be written as infinite union of disjoint point closures.

As every pair of non-empty  $at(\rho)$  - open sets intersect,  $at(\rho)$  cannot be  $S_{2\frac{1}{2}}$ . So the restriction  $L_\rho$  is well defined for characterizing minimal  $S_{2\frac{1}{2}}$  spaces.

**Lemma 3.5:** A space  $(X, \mathfrak{T})$  is  $S_{2\frac{1}{2}}$  if and only if it is  $S_1$  and  $adh_H N_{\mathfrak{T}}(x) = cl\{x\}$  for all  $x \in X$ .

**Proof:** Let  $(X, \mathfrak{T})$  be an  $S_{2\frac{1}{2}}$  space. Hence it is  $S_1$  also. Let  $x \in X$  and by Lemma 1.7 in an  $S_1$  space  $cl\{x\} \subset V$  for all  $V \in N_{\mathfrak{T}}(x)$ . So,  $cl\{x\} \subset cl_H V$  for all  $V \in N_{\mathfrak{T}}(x)$  and thus  $cl\{x\} \subset adh_H N_{\mathfrak{T}}(x)$ . Now let us suppose some  $y \in X$  be such that  $y \in adh_H N_{\mathfrak{T}}(x) \setminus cl\{x\}$ . Now  $y \in adh_H N_{\mathfrak{T}}(x)$  implies that  $y \in cl_H(V)$  for all  $V \in N_{\mathfrak{T}}(x)$ . Since  $V$  is an open set so it implies that for all  $U \in N_{\mathfrak{T}}(y)$ ,  $cl(U) \cap cl(V) \neq \emptyset$  for all  $V \in N_{\mathfrak{T}}(x)$ . As  $cl\{x\} \neq cl\{y\}$  and the space is  $S_{2\frac{1}{2}}$ , it is not possible. Thus it follows that  $adh_H N_{\mathfrak{T}}(x) = cl\{x\}$  for all  $x \in X$  as  $x$  was chosen arbitrarily. Conversely, suppose  $(X, \mathfrak{T})$  is  $S_1$  and  $adh_H N_{\mathfrak{T}}(x) = cl\{x\}$  for all  $x \in X$ . Let  $x$  and  $y$  be such that  $\langle x \rangle \neq \langle y \rangle$ . Then by Lemma 1.9 we have  $cl\{x\} \neq cl\{y\}$ . Since space  $X$  is  $S_1$  by Lemma 1.7  $cl\{x\} \cap cl\{y\} = \emptyset$ . Thus  $adh_H N_{\mathfrak{T}}(x) \cap adh_H N_{\mathfrak{T}}(y) = \emptyset$  which implies  $cl_H\{x\} \cap cl_H\{y\} = \emptyset$ . So by Lemma 1.8 the space  $(X, \mathfrak{T})$  is  $S_{2\frac{1}{2}}$ .

**Theorem 3.6:** Let  $\mathfrak{T} \in L_\rho$  be  $S_{2\frac{1}{2}}$ . Then the topology  $\mathfrak{T}$  is Minimal  $S_{2\frac{1}{2}}$  if and only if given any  $\mathfrak{T}$ -open filter  $F$  on  $X$  such that  $adh_{H(\mathfrak{T})} F = cl_{\mathfrak{T}}\{x\}$  for some  $x \in X$ , is convergent (necessarily to every point of  $cl_{\mathfrak{T}}\{x\}$ ).

**Proof:** Suppose  $\mathfrak{T}$  is Minimal  $S_{2\frac{1}{2}}$  and  $F$  is a  $\mathfrak{T}$ -open filter on  $X$  such that  $adh_{H(\mathfrak{T})} F = cl_{\mathfrak{T}}\{x\}$  for some  $x \in X$  and which is not convergent to some point  $z \in cl_{\mathfrak{T}}\{x\}$ . Since  $\mathfrak{T}$  is  $S_{2\frac{1}{2}}$  and hence  $S_1$  so  $cl_{\mathfrak{T}}\{x\} = cl_{\mathfrak{T}}\{z\}$ . So we can assume that  $F$  is not convergent to the point  $x$ .

Let us consider the topology  $\delta$  generated by the neighborhood base,

$$U^\delta(y) = \begin{cases} U, & \text{where } U \in N_\tau(y), y \neq x \\ U \cup F, & \text{where } U \in N_\tau(y) \text{ and } F \in F, y = x \end{cases}$$

Then  $\delta$  is a topology which is strictly coarser than  $\mathfrak{T}$ . We shall prove that  $\delta$  is an  $S_{2\frac{1}{2}}$  in  $L_\rho$ .

(1)  $\delta \in L_\rho$ : Since  $adh_{H(\mathfrak{T})} F = cl_{\mathfrak{T}}\{x\}$  and  $y \notin cl_{\mathfrak{T}}\{x\}$  then  $y \notin adh_{H(\mathfrak{T})} F$ . So there exists a  $V \in N_{\mathfrak{T}}(y)$  and  $F \in F$  such that  $cl(V) \cap cl(F) = \emptyset$ . Since  $cl_{\mathfrak{T}}\{y\} \subseteq cl(V)$  thus  $F \subseteq X \setminus cl_{\mathfrak{T}}\{y\}$ . Therefore  $X \setminus cl_{\mathfrak{T}}\{y\} \in F$  and hence  $N_{at(\rho)}(x) \subseteq F$ . Also, if  $N_{at(\rho)}(x) \subseteq F$  then  $N_{at(\rho)}(x) \subseteq N_\delta(x)$  for all  $x \in X$  which implies  $\delta \in L_\rho$ .

(2)  $\delta$  is  $S_{2\frac{1}{2}}$ : Now let  $y, z \in X$  be distinct points from given  $x$  such that  $y \notin cl_{\delta}\{z\}$ . Thus  $y \notin cl_{\mathfrak{T}}\{z\}$  and since  $\mathfrak{T}$  is  $S_{2\frac{1}{2}}$  there exists  $\mathfrak{T}$ -open sets  $U_1 \in N_{\mathfrak{T}}(y)$  and  $V_1 \in N_{\mathfrak{T}}(z)$  such that,  $cl_{\mathfrak{T}}(U_1) \cap cl_{\mathfrak{T}}(V_1) = \emptyset$ .

**Case I:** If  $x \notin cl_{\mathfrak{T}}\{y\}$  and  $x \notin cl_{\mathfrak{T}}\{z\}$  then  $y, z \notin cl_{\mathfrak{T}}\{x\}$ . Then  $y, z \notin adh_{H(\mathfrak{T})}F$  and so there exists  $F_1, F_2 \in F$  and  $U_2 \in N_{\mathfrak{T}}(y)$  and  $V_2 \in N_{\mathfrak{T}}(z)$  such that,  $cl_{\mathfrak{T}}(U_2) \cap cl_{\mathfrak{T}}(F_1) = \emptyset$  and  $cl_{\mathfrak{T}}(V_2) \cap cl_{\mathfrak{T}}(F_2) = \emptyset$ . Now since  $cl_{\mathfrak{T}}\{x\} \neq cl_{\mathfrak{T}}\{y\}$  and  $cl_{\mathfrak{T}}\{x\} \neq cl_{\mathfrak{T}}\{z\}$  and  $\mathfrak{T}$  is  $S_{2\frac{1}{2}}$  there exists  $U_3 \in N_{\mathfrak{T}}(y), V_3 \in N_{\mathfrak{T}}(z)$  and  $W_1, W_2 \in N_{\mathfrak{T}}(x)$  such that,  $cl_{\mathfrak{T}}(U_3) \cap cl_{\mathfrak{T}}(W_1) = \emptyset$  and  $cl_{\mathfrak{T}}(V_3) \cap cl_{\mathfrak{T}}(W_2) = \emptyset$ . Then  $O_y = U_1 \cap U_2 \cap U_3$  and  $O_z = V_1 \cap V_2 \cap V_3$  are  $\delta$ -open sets such that  $cl_{\mathfrak{T}}(O_y) \cap cl_{\mathfrak{T}}(O_z) = \emptyset$ . Since  $cl_{\delta}(O_y) = cl_{\mathfrak{T}}(O_y)$  and  $cl_{\delta}(O_z) = cl_{\mathfrak{T}}(O_z)$  we have  $cl_{\delta}(O_y) \cap cl_{\delta}(O_z) = \emptyset$ .

**Case II:** If  $x \in cl_{\mathfrak{T}}\{y\}$  then  $cl_{\mathfrak{T}}\{x\} = cl_{\mathfrak{T}}\{y\} = adh_{H(\mathfrak{T})}F$ . Since  $z \notin cl_{\mathfrak{T}}\{y\}$  there exist  $V_4 \in N_{\mathfrak{T}}(z)$  and  $F_3 \in F$  such that  $cl_{\mathfrak{T}}(V_4) \cap cl_{\mathfrak{T}}(F_3) = \emptyset$ . Since  $cl_{\mathfrak{T}}\{x\} \neq cl_{\mathfrak{T}}\{z\}$   $W_3 \in N_{\mathfrak{T}}(x)$  and  $V_5 \in N_{\mathfrak{T}}(z)$  such that,  $cl_{\mathfrak{T}}(W_3) \cap cl_{\mathfrak{T}}(V_5) = \emptyset$ . Then  $P_y = W_3 \cup F_3$  and  $P_z = V_4 \cap V_5$  are  $\delta$ -open sets such that  $cl_{\mathfrak{T}}(P_y) \cap cl_{\mathfrak{T}}(P_z) = \emptyset$ . Since  $cl_{\delta}(P_y) = cl_{\mathfrak{T}}(P_y)$  and  $cl_{\delta}(P_z) = cl_{\mathfrak{T}}(P_z)$  we have  $cl_{\delta}(P_y) \cap cl_{\delta}(P_z) = \emptyset$ .

Thus  $\delta$  is an  $S_{2\frac{1}{2}}$  topology coarser than  $\mathfrak{T}$  in  $L_p$  which is a contradiction and the result follows.

Conversely, let every  $\mathfrak{T}$ -open filter  $F$  on  $X$  such that  $adh_{H(\mathfrak{T})}F = cl_{\mathfrak{T}}\{x\}$  for some  $x \in X$ , is convergent. Let  $\mathfrak{T}^* \in L_p$  be  $S_{2\frac{1}{2}}$  such that  $\mathfrak{T}^* \leq \mathfrak{T}$ . Let  $x \in U$  for some set  $U \in \mathfrak{T}$ . Then  $adh_{H(\mathfrak{T}^*)}N_{\mathfrak{T}^*}(x) = cl_{\mathfrak{T}}\{x\}$  and so  $\mathfrak{T}$ -open filter  $N_{\mathfrak{T}^*}(x)$  is  $\mathfrak{T}$ -convergent to  $x$ . So  $N_{\mathfrak{T}}(x) \subseteq N_{\mathfrak{T}^*}(x)$  and hence  $U \in \mathfrak{T}^*$ . As  $U$  was chosen arbitrarily we have  $\mathfrak{T}^* = \mathfrak{T}$  and the result follows.

As a consequence of Theorem 3.6 we obtain that every Minimal  $S_{2\frac{1}{2}}$  space is  $U(i)$ .

**Corollary 3.7:** If  $\mathfrak{T} \in L_p$  is Minimal  $S_{2\frac{1}{2}}$  then it is  $U(i)$ .

**Proof:** Suppose there exists an open filter  $F$  such that  $adh_{H(\mathfrak{T})}F = \emptyset$ . Then for each  $x \in X$  there exists a  $V \in N_{\mathfrak{T}}(x)$  and  $F \in F$  such that  $cl_{\mathfrak{T}}(V) \cap cl_{\mathfrak{T}}(F) = \emptyset$ . On the other hand since  $cl_{\mathfrak{T}}\{x\} \subseteq V$  then  $X \setminus cl_{\mathfrak{T}}\{x\} \in F$ . Thus  $N_{at(\rho)}(x) \subseteq F$  and  $N_{\mathfrak{T}}(x) \not\subseteq F$  for each  $x \in X$ . Now fix  $x \in X$  and  $\delta$  be defined as in Theorem 3.6 above. Then  $\delta \in L_p$  and  $\delta \leq \mathfrak{T}$ . For proving  $\delta$  is  $S_{2\frac{1}{2}}$  we show that  $F$  contains an open filter  $F^*$  such that  $adh_{H(\mathfrak{T})}F^* = cl_{\mathfrak{T}}\{x\}$ .

Let  $F^* = \{F \in F : cl_{\mathfrak{T}}(U) \cap cl_{\mathfrak{T}}(F) \neq \emptyset \text{ for all } U \in N_{\mathfrak{T}}(x)\}$ . Then  $F^*$  is a proper sub-filter of  $F$  and  $cl_{\mathfrak{T}}\{x\} \subseteq adh_{H(\mathfrak{T})}F^*$ . Now let  $y \notin cl_{\mathfrak{T}}\{x\}$  and as  $\mathfrak{T}$  is  $S_{2\frac{1}{2}}$  there exists  $\mathfrak{T}$ -open sets  $U_1 \in N_{\mathfrak{T}}(x)$  and  $V_1 \in N_{\mathfrak{T}}(y)$  such that,  $cl_{\mathfrak{T}}(U_1) \cap cl_{\mathfrak{T}}(V_1) = \emptyset$ . Since  $y \notin adh_{H(\mathfrak{T})}F$  there exist  $W_1 \in N_{\mathfrak{T}}(y)$   $F_1 \in F$  such that,  $cl_{\mathfrak{T}}(W_1) \cap cl_{\mathfrak{T}}(F_1) = \emptyset$ . Then  $O = W_1 \cap V_1$  and  $G = U_1 \cup F_1$  are such that  $cl_{\mathfrak{T}}(O) \cap cl_{\mathfrak{T}}(G) = \emptyset$  where  $O \in N_{\mathfrak{T}}(y)$  and  $G \in F^*$ . So  $y \notin adh_{H(\mathfrak{T})}F^*$  and hence  $adh_{H(\mathfrak{T})}F^* = cl_{\mathfrak{T}}\{x\}$ . Then by similar steps as in Theorem 3.6  $\delta$  is  $S_{2\frac{1}{2}}$  which implies that  $\mathfrak{T}$  is not Minimal  $S_{2\frac{1}{2}}$  which is a contradiction and the result follows.

## REFERENCES

- [1] M. P. Berri, J. R. Porter, R. M. Stephenson, A survey of minimal topological spaces, Proc. Kanpur Topology Conference 1968, Academic Press (1970), 93-114.
- [2] L. L. Herrington, Characterizations of Urysohn-closed spaces, Proc. Amer. Math. Soc., **53** (1976), 435-439.
- [3] N.S. Noorie and Anakh Singh, On Hyperclosed Sets, Int. Journal of Pure and Applied Mathematics, **102(1)** (2015), 39-50.
- [4] C. T. Scarborough, Minimal Urysohn spaces, Pacific J. Math., **27** (1968), 611- 617.
- [5] M. L. Colasante and D. Van der Zypen, Minimal  $R_1$ , minimal regular and minimal presober topologies, Revista Notas de Matemática, **5(1) (275)**, (2009), 73-84.
- [6] J. K. Kohli and A. K. Das, A class of spaces containing all generalized absolutely closed (almost compact) spaces, Applied General Topology, **7(2)**(2006), 233-244.
- [7] D. Janković, On some separation axioms and  $\theta$ -closure, Mat. Vesnik, **32(4) (17)** (1980), 439-449.